lyn, NY 11218 New York,	attan Street 601 NY 10019 Threewell	M	Manhattan 225 E 70th Street Suite 1E New York, NY 10021	Queens 64-05 Yellowstone Blvd CF104 Forest Hills, NY 11375	Manhattan 225 East 70th St New York, NY 1
tt Shore Road 165 North Village Avenue 555 Ta nite 201 Suite 133 3rd	d/Terrytown I N F U S I O N axter Road d Floor Office: 212-803-3339 Fax: 646-768-8600 rd, NY 10523	Mission Medical	Holbrook/Ronkonkoma 233 Union Ave Suite 207 Holbrook, NY 11741	Scarsdale 495 Central Park Avenue Suite 205 Scarsdale, NY 10583	5 Towns 141 Washington Av Cedarhurst, NY 1
INFUSION OR			Long Beach 917 Beech Street Long Beach, NY 11561	Riverhead 1228 E Main Street Suite A	
	$\overline{{ m IS}}_{ m (INFLIXIMAB-abda)}$ Date:			Riverhead, NY 11901	
Name	PATIENT INFORM	MATION			
Name: Allergies:	DOB: Date of Refe	rral:			
	REFERRAL STATU	S			
	New Referral ☐ Dose or Frequency Char	nge 🗆 Order	Renewal		
	INFUSION OFFICE PREFERE	NCES (Optional)			
Preferred Location*:					
	DIAGNOSIS AND ICD				
☐ Moderate to Severe Ulcerati					
☐ Moderate to Severe Crohn's					
☐ Rheumatoid Arthritis	ICD 10 Code: ICD 10 Code:				
☐ Ankylosing Spondylitis	ICD 10 Code:				
☐ Psoriatic Arthritis	ICD 10 Code:				
☐ Plaque Psoriasis ☐ Other:	ICD 10 Code:				
Other.	ICD10 Code.				
	required documen				
☐ This signed order form by the	•		Progress notes		
□ Patient demographics AND insurance information□ Hepatitis B Test Results: HBsAg, Total HepB Core Antibody			☐ Labs and Tests supporting primary diagnosis☐ TB Test Results		
List Tried & Failed Therapies, in	· , ,		Results		
1)	retaining duration of treatment.				
2)					
3)					
	MEDICATION ORE	DERS			
	☐ Renflexis 5mg/kg IV at week 0, 2, 6, th	nen every 8 weeks t	hereafter		
Initial Dosing					
	☐ Renflexis 5mg/kg IV every 8 weeks				
Initial Dosing Maintenance Dosing Alternative Dosing		weeks			
Maintenance Dosing Alternative Dosing Patient Weight=	☐ Renflexis 5mg/kg IV every 8 weeks ☐ Renflexis IV every kg	weeks			
Maintenance Dosing Alternative Dosing Patient Weight=	☐ Renflexis 5mg/kg IV every 8 weeks ☐ Renflexis IV every _				
Maintenance Dosing Alternative Dosing Patient Weight= Refills:	Renflexis 5mg/kg IV every 8 weeks Renflexis IV every Weeks kg 6 months X 1 year dose	es NS			
Maintenance Dosing Alternative Dosing Patient Weight= Refills: \(\sum X \)	Renflexis 5mg/kg IV every 8 weeks Renflexis IV every IV every kg 6 months X 1 year dose PREMEDICATION prior to Remicade infusion	es FREQUENCY			
Maintenance Dosing Alternative Dosing Patient Weight= Refills:	Renflexis 5mg/kg IV every 8 weeks Renflexis IV every kg 6 months X 1 year dose PREMEDICATION prior to Remicade infusion prior to Remicade infusion	FREQUENCY Week 2, 6, the	en every 8 weeks		
Maintenance Dosing Alternative Dosing Patient Weight= Refills:	Renflexis 5mg/kg IV every 8 weeks Renflexis IV every IV every kg 6 months X 1 year dose PREMEDICATION prior to Remicade infusion	FREQUENCY Week 2, 6, th Every 6 weeks	3		
Maintenance Dosing Alternative Dosing Patient Weight= Refills: Acetaminophen 650mg PO Diphenhydramine 25mg PO Methylprednisolone 40mg S Other:	Renflexis 5mg/kg IV every 8 weeks Renflexis IV every IV every kg 6 months X 1 year dose PREMEDICATION prior to Remicade infusion prior to Remicade infusion Slow IV Push PRN infusion reaction	FREQUENCY Week 2, 6, th Every 6 weeks Every 8 weeks	· ·	ad medically no	
Maintenance Dosing Alternative Dosing Patient Weight= Refills:	Renflexis 5mg/kg IV every 8 weeks Renflexis IV every kg 6 months X 1 year dose PREMEDICATION prior to Remicade infusion prior to Remicade infusion	FREQUENCY FREQUENCY Week 2, 6, th Every 6 weeks Every 8 weeks	· ·	ed medically ne	cessary.
Maintenance Dosing Alternative Dosing Patient Weight= Refills:	Renflexis 5mg/kg IV every 8 weeks Renflexis IV every Vevery Vever Vevery Vever Vevery Vever Vevery Vever Vever Vevery Vevery Vevery Vever	FREQUENCY Week 2, 6, th Every 6 weeks Every 8 weeks ropriate rescue medication.	· ·	ed medically ne	ecessary.
Maintenance Dosing Alternative Dosing Patient Weight= Refills:	Renflexis 5mg/kg IV every 8 weeks Renflexis IV every kg 6 months X 1 year dose PREMEDICATION prior to Remicade infusion prior to Remicade infusion Slow IV Push PRN infusion reaction on occurs, the on-call physician will order appeaducing the rate of infusion or discontinuing the PRESCRIBER INFORM	FREQUENCY FREQUENCY Week 2, 6, th Every 6 weeks Every 8 weeks ropriate rescue medication.	dications as deeme	ed medically ne	ccessary.
Maintenance Dosing Alternative Dosing Patient Weight= Refills:	Renflexis 5mg/kg IV every 8 weeks Renflexis IV every kg 6 months X 1 year dose PREMEDICATION prior to Remicade infusion prior to Remicade infusion Oprior to Remicade infusion Slow IV Push PRN infusion reaction on occurs, the on-call physician will order appeaducing the rate of infusion or discontinuing the	FREQUENCY Week 2, 6, the Every 6 weeks Every 8 weeks ropriate rescue medication. ATION	dications as deeme	ed medically ne	ecessary.
Maintenance Dosing Alternative Dosing Patient Weight= Refills:	Renflexis 5mg/kg IV every 8 weeks Renflexis IV every kg 6 months X 1 year dose PREMEDICATION prior to Remicade infusion prior to Remicade infusion Slow IV Push PRN infusion reaction on occurs, the on-call physician will order appeaducing the rate of infusion or discontinuing the PRESCRIBER INFORM	FREQUENCY FREQUENCY Week 2, 6, th Every 6 weeks Every 8 weeks ropriate rescue medication.	dications as deeme	ed medically ne	ecessary.
Maintenance Dosing Alternative Dosing Patient Weight= Refills:	Renflexis 5mg/kg IV every 8 weeks Renflexis IV every kg 6 months X 1 year dose PREMEDICATION prior to Remicade infusion prior to Remicade infusion Slow IV Push PRN infusion reaction on occurs, the on-call physician will order appeducing the rate of infusion or discontinuing the PRESCRIBER INFORM Office Fax:	FREQUENCY Week 2, 6, the Every 6 weeks Every 8 weeks ropriate rescue medication. ATION	dications as deeme	ed medically ne	ecessary.
Maintenance Dosing Alternative Dosing Patient Weight= Refills:	Renflexis 5mg/kg IV every 8 weeks Renflexis IV every kg 6 months X 1 year dose PREMEDICATION prior to Remicade infusion prior to Remicade infusion Slow IV Push PRN infusion reaction on occurs, the on-call physician will order appeducing the rate of infusion or discontinuing the PRESCRIBER INFORM Office Fax:	FREQUENCY Week 2, 6, the Every 6 weeks Every 8 weeks ropriate rescue medication. ATION	dications as deeme	ed medically ne	ecessary.

Provider Phone Fax