

(rituximab)

# RITUXAN infusion orders

Date: \_\_\_\_\_

Patient Name \_\_\_\_\_ DOB \_\_\_\_\_

Phone \_\_\_\_\_ ☐ Allergies M ☐ F ☐

NPI \_\_\_\_\_ Tax ID \_\_\_\_\_

Insurance Carrier (primary) \_\_\_\_\_

Insurance Carrier (secondary) \_\_\_\_\_

## REFERRAL STATUS

- ☐ New Prescription  
☐ Order Renewal  
☐ Dose or Frequency Change  
☐ Discontinuation

## DIAGNOSIS Please provide ICD-10 code

☐ \_\_\_\_\_ Rheumatoid Arthritis

☐ \_\_\_\_\_ Granulomatosis w/Polyangitis  
(wegener's granulomatosis GPA)

☐ \_\_\_\_\_ Microscopic Polyangitis

☐ \_\_\_\_\_  
(other)

## PRE-MEDICATION

☐ Tylenol 1000mg PO

☐ Diphenhydramine 25mg PO

☐ Cetirizine 10mg PO

☐ \_\_\_\_\_

☐ Solu-Medrol 125mg IVP

☐ Solu-Cortef 100mg IVP

☐ Diphenhydramine 25mg IVP

☐ \_\_\_\_\_

## RITUXAN ORDERS

### DOSAGE

☐ 1000mg

☐ 375mg/m<sup>2</sup>

☐ Other \_\_\_\_\_

### FREQUENCY

☐ initial dose (0) followed by 2nd dose on day 15 (induction for RA diagnosis)

☐ single dose

☐ every week for 4 weeks total

☐ \_\_\_\_\_  
(other frequency)

### PATIENT WEIGHT

\_\_\_\_\_ lbs.

\_\_\_\_\_ kg

☐ Total dosages \_\_\_\_\_

☐ Refills \_\_\_\_\_

## NOTES

## ORDERING PROVIDER

Signature **X** \_\_\_\_\_ Date \_\_\_\_\_

Provider \_\_\_\_\_ Phone \_\_\_\_\_ Fax \_\_\_\_\_