ough Park   Crown Heights   S55 Lefferts Avenue   S7W 57Street   Suite 601   New York, NY 1019	ewell 🚜	2251	anhattan E 70th Street Suite 1E ork, NY 10021	Queens 64-05 Yellowstone Blvd CF104 Forest Hills, NY 11375	Manhattan 225 East 70th Str New York, NY 10
thasset	I O N Mission N	)	lbrook/ Ronkonkoma 233 Union Ave Suite 207 Holbrook, NY 11741	495 Central Park Avenue Suite 205 Scarsdale, NY 10583	5 Towns 141 Washington Ave Cedarhurst, NY 115
TEDE77 A		917 1	ong Beach Beech Street each, NY 11561	Riverhead  1228 E Main Street Suite A	
TEPEZZA INFUSION ORDE				Riverhead, NY 11901	
<b>Note:</b> This form is being provided as a guide. Prescribe acilities prefer to use their own infusion order form. Cl					der.
•	TENT INFORMATION		0 7		
Name:	DOB:	Sex: M	F \	Weight: kilo□ II	b□
Phone number:	Email:	'	<u>'</u>		
Allergies:	ICD-10 code:				
Is the patient diabetic? Yes ☐ No ☐	Does the patient	have a history	of IBD?	Yes□ No□	
Emergency contact name:	Phone number:	,			
Please attach: 1. List of current n 3. Clinical progress notes and history and					
PHYS	SICIAN INFORMAT	ΓΙΟΝ			
Prescribing Physician's Name:	Practice Name:				
Phone Number:	Fax Number:				
Email:	Office Contact:				
Co-managing Physician Name:	Phone Number/E	Email:			
Medication: TEPEZZA (teprotumumab-trbw)  Dose: Infusion 1: mg (10 mg/kg) Infusions 2 to 8: (20 mg/kg)  Duration: Administer the first 2 infusions over 90 minutes. note below for additional information).  Saline bag: Administer via an infusion bag containing 0.9%	Subsequent infusions may k				
Medication: TEPEZZA (teprotumumab-trbw) Dose: Infusion 1: mg (10 mg/kg) Infusions 2 to 8: (20 mg/kg) Duration: Administer the first 2 infusions over 90 minutes. note below for additional information). Saline bag: Administer via an infusion bag containing 0.9% For doses 1800 mg, use a 250-mL bag.	3) Subsequent infusions may b 5 Sodium Chloride Solution,	ı, USP. For dose			
Medication: TEPEZZA (teprotumumab-trbw)  Dose: Infusion 1: mg (10 mg/kg) Infusions 2 to 8: (20 mg/kg)  Duration: Administer the first 2 infusions over 90 minutes. note below for additional information).  Saline bag: Administer via an infusion bag containing 0.9%	Subsequent infusions may be Sodium Chloride Solution,  Pretreatment medi	, USP. For dose	es <1800 m	g, use a100-mL	bag.
Medication: TEPEZZA (teprotumumab-trbw) Dose: Infusion 1: mg (10 mg/kg) Infusions 2 to 8: (20 mg/kg) Duration: Administer the first 2 infusions over 90 minutes. note below for additional information). Saline bag: Administer via an infusion bag containing 0.9% For doses 1800 mg, use a 250-mL bag.	3) Subsequent infusions may b 5 Sodium Chloride Solution,	ications:  pes not require a syprotocol. If the partition for subsequents	es <1800 mg	g, use a100-mL	bag.
Medication: TEPEZZA (teprotumumab-trbw) Dose: Infusion 1: mg (10 mg/kg) Infusions 2 to 8: (20 mg/kg) Duration: Administer the first 2 infusions over 90 minutes. note below for additional information). Saline bag: Administer via an infusion bag containing 0.9% For doses 1800 mg, use a 250-mL bag.  Schedule: Q3 weeks, 8 infusions total	Subsequent infusions may be Sodium Chloride Solution, Pretreatment medi Note: TEPEZZA do follow your facility consider premedic additional informatics.	ications:  pes not require a syprotocol. If the partition for subsequition).	specific proto patient experi lent doses (se	g, use a100-mL  ocol for premedica iences an infusion ee note below for	tions;
Medication: TEPEZZA (teprotumumab-trbw) Dose: Infusion 1: mg (10 mg/kg) Infusions 2 to 8: (20 mg/kg) Duration: Administer the first 2 infusions over 90 minutes. note below for additional information). Saline bag: Administer via an infusion bag containing 0.9% For doses 1800 mg, use a 250-mL bag.  Schedule: Q3 weeks, 8 infusions total  Preferred start date:   If an infusion reaction occurs, interrupt or slow the rate of infusion to 90 minutes and consider premedicating with  Follow your facility protocol and notify the prescriber. For flush solution, declotting, and/or dressing changes.	Subsequent infusions may be Sodium Chloride Solution, Pretreatment medical Note: TEPEZZA do follow your facility consider premedical additional information and use appropriate an antihistamine, antipyretic, a	ications:  pes not require a syprotocol. If the pation for subsequition).	specific proto patient experi lent doses (se	g, use a100-mL ocol for premedicationces an infusion see note below for obsequent infusions	ations; reaction,
Medication: TEPEZZA (teprotumumab-trbw) Dose: Infusion 1: mg (10 mg/kg) Infusions 2 to 8: (20 mg/kg) Duration: Administer the first 2 infusions over 90 minutes. note below for additional information). Saline bag: Administer via an infusion bag containing 0.9% For doses 1800 mg, use a 250-mL bag.  Schedule: Q3 weeks, 8 infusions total  Preferred start date:   If an infusion reaction occurs, interrupt or slow the rate of infusion to 90 minutes and consider premedicating with flush solution, declotting, and/or dressing changes.  Share post-infusion chart notes with the prescriber.	Subsequent infusions may be Sodium Chloride Solution, Pretreatment medical Note: TEPEZZA do follow your facility consider premedical additional information and use appropriate an antihistamine, antipyretic, a	ications:  pes not require a syprotocol. If the pation for subsequition).	specific proto patient experi lent doses (se	g, use a100-mL ocol for premedicationces an infusion see note below for obsequent infusions	ations; reaction,
Medication: TEPEZZA (teprotumumab-trbw) Dose: Infusion 1: mg (10 mg/kg) Infusions 2 to 8: (20 mg/kg) Duration: Administer the first 2 infusions over 90 minutes. note below for additional information). Saline bag: Administer via an infusion bag containing 0.9% For doses 1800 mg, use a 250-mL bag.  Schedule: Q3 weeks, 8 infusions total  Preferred start date:  Notes:  If an infusion reaction occurs, interrupt or slow the rate of infusion to 90 minutes and consider premedicating with  Follow your facility protocol and notify the prescriber. For flush solution, declotting, and/or dressing changes.  Share post-infusion chart notes with the prescriber.  Other notes:	Subsequent infusions may be Sodium Chloride Solution, Pretreatment medical Note: TEPEZZA do follow your facility consider premedical additional information of infusion and use appropriate an antihistamine, antipyretic, a	ications:  pes not require a syprotocol. If the pation for subsequition).	specific proto patient experi lent doses (se	g, use a100-mL ocol for premedicationces an infusion see note below for obsequent infusions	ations; reaction,
Medication: TEPEZZA (teprotumumab-trbw) Dose: Infusion 1: mg (10 mg/kg) Infusions 2 to 8: (20 mg/kg) Duration: Administer the first 2 infusions over 90 minutes. note below for additional information). Saline bag: Administer via an infusion bag containing 0.9% For doses 1800 mg, use a 250-mL bag.  Schedule: Q3 weeks, 8 infusions total  Preferred start date:  Notes:  If an infusion reaction occurs, interrupt or slow the rate of infusion to 90 minutes and consider premedicating with flush solution, declotting, and/or dressing changes.  Share post-infusion chart notes with the prescriber.  Other notes:	Subsequent infusions may be Sodium Chloride Solution, Pretreatment medical Note: TEPEZZA do follow your facility consider premedical additional information and use appropriate an antihistamine, antipyretic, a	ications:  pes not require a syprotocol. If the pation for subsequition).	specific proto patient experi lent doses (se	g, use a100-mL ocol for premedicationces an infusion see note below for obsequent infusions	ations; reaction,
Medication: TEPEZZA (teprotumumab-trbw) Dose: Infusion 1: mg (10 mg/kg) Infusions 2 to 8: (20 mg/kg) Duration: Administer the first 2 infusions over 90 minutes. note below for additional information). Saline bag: Administer via an infusion bag containing 0.9% For doses 1800 mg, use a 250-mL bag.  Schedule: Q3 weeks, 8 infusions total  Preferred start date:	Subsequent infusions may be Sodium Chloride Solution, Pretreatment medical Note: TEPEZZA do follow your facility consider premedical additional information of infusion and use appropriate an antihistamine, antipyretic, a	ications:  pes not require a syprotocol. If the pation for subsequition).	specific proto patient experi lent doses (se	g, use a100-mL ocol for premedicationces an infusion see note below for obsequent infusions	ations; reaction,
Medication: TEPEZZA (teprotumumab-trbw) Dose: Infusion 1: mg (10 mg/kg) Infusions 2 to 8: (20 mg/kg) Duration: Administer the first 2 infusions over 90 minutes. note below for additional information). Saline bag: Administer via an infusion bag containing 0.9% For doses 1800 mg, use a 250-mL bag.  Schedule: Q3 weeks, 8 infusions total  Preferred start date:	Subsequent infusions may be Sodium Chloride Solution, Pretreatment medical Note: TEPEZZA do follow your facility consider premedical additional information of infusion and use appropriate an antihistamine, antipyretic, a collow facility policies and/or presented to the property of the	ications:  pes not require a syprotocol. If the pation for subsequation).  medical manage and/or corticoster rotocols for vascu	specific proto patient experi lent doses (se	g, use a100-mL ocol for premedicationces an infusion see note below for obsequent infusions	ations; reaction,
Medication: TEPEZZA (teprotumumab-trbw)  Dose: Infusion 1: mg (10 mg/kg) Infusions 2 to 8: (20 mg/kg)  Duration: Administer the first 2 infusions over 90 minutes. note below for additional information).  Saline bag: Administer via an infusion bag containing 0.9% For doses 1800 mg, use a 250-mL bag.  Schedule: Q3 weeks, 8 infusions total  Preferred start date:  If an infusion reaction occurs, interrupt or slow the rate of infusion to 90 minutes and consider premedicating with glush solution, declotting, and/or dressing changes.    Share post-infusion chart notes with the prescriber. Other notes:    Other notes:   infusion(s)	Subsequent infusions may be Sodium Chloride Solution, Pretreatment medical Note: TEPEZZA do follow your facility consider premedical additional information of infusion and use appropriate an antihistamine, antipyretic, a collow facility policies and/or presented to the property of the	ications:  pes not require a syprotocol. If the pation for subsequation).  medical manage and/or corticoster rotocols for vascuibing Information.	specific proto patient experi lent doses (se	g, use a100-mL ocol for premedicationces an infusion see note below for obsequent infusions	tions; reaction,
Medication: TEPEZZA (teprotumumab-trbw)  Dose: Infusion 1: mg (10 mg/kg) Infusions 2 to 8: (20 mg/kg)  Duration: Administer the first 2 infusions over 90 minutes. note below for additional information).  Saline bag: Administer via an infusion bag containing 0.9% For doses 1800 mg, use a 250-mL bag.  Schedule: Q3 weeks, 8 infusions total  Preferred start date:  If an infusion reaction occurs, interrupt or slow the rate of infusion to 90 minutes and consider premedicating with glush solution, declotting, and/or dressing changes.    Share post-infusion chart notes with the prescriber. Other notes:    Other notes:   infusion(s)	Subsequent infusions may be Sodium Chloride Solution, Pretreatment medical Note: TEPEZZA do follow your facility consider premedical additional information and use appropriate an antihistamine, antipyretic, a collow facility policies and/or prescription.  LAB ORDERS  accompanying Full Prescriptions in Substitution of the Companying Full Prescription in Substitution in	ications:  pes not require a syprotocol. If the pation for subsequation).  medical manage and/or corticoster rotocols for vascuibing Information.	specific proto patient experi lent doses (se	g, use a100-mL ocol for premedicationces an infusion see note below for obsequent infusions	tions; reaction,
Medication: TEPEZZA (teprotumumab-trbw)  Dose: Infusion 1: mg (10 mg/kg) Infusions 2 to 8: (20 mg/kg)  Duration: Administer the first 2 infusions over 90 minutes. note below for additional information).  Saline bag: Administer via an infusion bag containing 0.9% For doses 1800 mg, use a 250-mL bag.  Schedule: Q3 weeks, 8 infusions total  Preferred start date:  If an infusion reaction occurs, interrupt or slow the rate of infusion to 90 minutes and consider premedicating with glush solution, declotting, and/or dressing changes.    Share post-infusion chart notes with the prescriber. Other notes:    Other notes:   infusion(s)	Subsequent infusions may be Sodium Chloride Solution,  Pretreatment medical Note: TEPEZZA do follow your facility consider premedical additional information and use appropriate an antihistamine, antipyretic, a collow facility policies and/or prediction of the solution o	ications:  pes not require a syprotocol. If the pation for subsequation).  medical manage and/or corticoster rotocols for vascustions for vascustions.	specific proto patient experi lent doses (se	g, use a100-mL ocol for premedicationces an infusion see note below for obsequent infusions	tions; reaction,
Medication: TEPEZZA (teprotumumab-trbw)  Dose: Infusion 1: mg (10 mg/kg) Infusions 2 to 8: (20 mg/kg)  Duration: Administer the first 2 infusions over 90 minutes. note below for additional information).  Saline bag: Administer via an infusion bag containing 0.9% For doses 1800 mg, use a 250-mL bag.  Schedule: Q3 weeks, 8 infusions total  Preferred start date:	Subsequent infusions may be Sodium Chloride Solution, Sodium Chloride Solution, Pretreatment medical Note: TEPEZZA do follow your facility consider premedical additional information of infusion and use appropriate an antihistamine, antipyretic, a collow facility policies and/or prediction of the properties of the pro	ications:  pes not require a syprotocol. If the pation for subsequation).  medical manage and/or corticoster rotocols for vascustions for vascustions.	specific proto patient experi lent doses (se	g, use a100-mL ocol for premedicationces an infusion see note below for obsequent infusions	ations; reaction,

Second Insurance

Policy #/ Group #