Chicago Illinois 4711 Golf Road Suite 900 Skokie, IL 60076





Canakinumab (Ilaris) Provider Order Form

Provider Order Form	Date:	
PATIENT	INFORMATION	
Name:	DOB: SEX: M \(\sigma \) F \(\sigma \)	
ICD-10 code (required):	ICD-10 description:	
□NKDA Allergies:	Weight lbs/kg:	
REFERR	AL STATUS	
□New Referral □Referral Renewal □Medication/Order C		er
	N INFORMATION	
Referral Coordinator Name:	Referral Coordinator Email:	
Ordering Provider:	Provider NPI:	
Referring Practice Name:	Phone: Fax:	
Practice Address:	City: State: Zip Code:	
Patient is required to stay for 30 minutes observation period Patient is NOT required to stay for observation time Other: SPECIAL INSTRUCTIONS NOTES/ADDITIONAL COMMENTS:	Canakinumab (Ilaris) For Stills Disease including Adult Onset Stills Disease and Systemic Juvenile Idiopathic Arthritis. □ 4mg/kg (with a max of 300mg) for patients with a body weight greater than or equal to 7.5kg subcutaneous every 4 weeks □ Other □ For Cryopyrin-Associated Periodic Syndromes (CAPS) □ 150mg for patients with body weight greater than 40kg subcutaneous every 8 weeks □ 2mg/kg for patients with body weight greater than or equal to 15kg and less than or equal to 40kg subcutaneous every 8 wks □ Other □ For Tumor Necrosis Factor Receptor Associated Periodic Syndrome, Hyperimmunoglobulin D Syndrome/Mevalonate Kinase Deficiency, Familial Mediterranean Fever Body weight less than or equal to 40kg □ 2mg/kg subcutaneous every 4 weeks □ 4mg/kg subcutaneous every 4 weeks - consider if clinical □ Other □ responsenot adequate. Body weight greater than 40kg □ 150mg subcutaneous every 4 weeks - consider if clinical response not adequate. Refills:□Zero / □ for 12 months / □ □ (if not indicated order will expire one year from date signed) □ Other □ □ Total Doses □ □ Refills □ □	
ORDERING PROVIDER Signature X	Date	
Provider	Phone Fax	