

- Borough Park**
1428 36th Street
Suite 107
Brooklyn, NY 11218
- Crown Heights**
555 Lefferts Avenue
Brooklyn, NY 11225
- Manhattan**
57W 57Street
Suite 601
New York, NY 10019
- Manhasset**
333 East Shore Road
Suite 201
Manhasset, NY 11030
- Rockville Centre**
165 North Village Avenue
Suite 133
Rockville Center, NY 11570
- Elmsford/Tarrytown**
555 Taxter Road
3rd Floor
Elmsford, NY 10523
- NYC Central Park West**
115 Central Park West
Suite 15
New York, NY 10023
- Woodbury**
75 Froehlich Farm
Woodbury, NY 11797
- Staten Island**
27 New Dorp Lane
Staten Island, NY 10306



- Manhattan**
225 E 70th Street
Suite 1E
New York, NY 10021
- Queens**
64-05 Yellowstone Blvd
CF104
Forest Hills, NY 11375
- Manhattan**
225 East 70th Street
New York, NY 10021
- Holbrook/Ronkonkoma**
233 Union Ave
Suite 207
Holbrook, NY 11741
- Scarsdale**
495 Central Park Avenue
Suite 205
Scarsdale, NY 10583
- 5 Towns**
141 Washington Avenue
Cedarhurst, NY 11559
- Long Beach**
917 Beech Street
Long Beach, NY 11561
- Riverhead**
1228 E Main Street
Suite A
Riverhead, NY 11901
- Bronx**
226 West 238th Street
Bronx, NY 10463

Provider Order Form

Iron (Feraheme/Injectafer/Venofer) Date: _____

PATIENT INFORMATION

Name:	DOB:
Allergies:	Date of Referral:

ICD-10 code (required): _____ ICD -10 description: _____

NKDA Allergies: _____ Weight lbs/kg: _____

Patient Status: New to Therapy Continuing Therapy Next Due Date (if applicable): _____

REFERRAL STATUS: New Prescription Order Renewal Does or Frequency Change Discontinuation

PROVIDER INFORMATION

Referral Coordinator Name: _____ Referral Coordinator Email: _____

Ordering Provider: _____ Provider NPI: _____

Referring Practice Name: _____ Phone: _____ Fax: _____

Practice Address: _____ City: _____ State: _____ Zip Code: _____

PREN-MEDICATION ORDERS

- acetaminophen (Tylenol) 500mg / 650mg / 1000mg PO
- cetirizine (Zyrtec) 10mg PO
- loratadine (Claritin) 10mg PO
- diphenhydramine (Benadryl) 25mg / 50mg PO / IV
- methylprednisolone (Solu-Medrol) 40mg / 125mg IV
- Other: _____
Dose: _____ Route: _____
Frequency: _____

SPECIAL INSTRUCTIONS

*Closely observe patients for signs and symptoms of hypersensitivity including monitoring of blood pressure and pulse during and after Feraheme administration for at least 30 minutes and until clinically stable following completion of each infusion.

Observe for signs and symptoms of hypersensitivity during and after Injectafer administration for at least 30 minutes and until clinically stable following completion of each administration.*Monitor patients for signs and symptoms of hypersensitivity during and after Venofer administration for at least 30 minutes and until clinically

THERAPY ADMINISTRATION

- Ferumoxylol (Feraheme) intravenous infusion
 - Dose & Frequency: initial 510mg infusion followed by a second 510mg infusion 3-8 days later
 - Diluten 50 - 200ml 0.9% sodium chloride or 5% dextrose solution (final concentration 2mg - 8mg per ml)
 - Infuse over at least 15 minutes
 - No refills Other
- Ferriccarboxymaltose (Injectafer) intravenous infusion
 - Dose & Frequency: Patients > 50kg: Two 750mg doses, 7 days apart / Patients < 50kg: Two 15mg/kg doses, 7 days apart
 - Diluten no more than 250ml 0.9% sodium chloride
 - Infuse over at least 15 minutes
 - No refills Other
- Ironsucrose(Venofer) intravenous infusion
 - Dose:
 - 100mg in 100ml 0.9% sodium chloride over 30 minutes
 - 200mg in 100ml 0.9% sodium chloride over 30minutes
 - 300mg in 250ml 0.9% sodium chloride over 1.5 hours
 - 400mg in 250ml 0.9% sodium chloride over 2.5 hours
 - _____
 - Frequency:
 - Once Every 2- 3 days x _____ doses
 - Daily x _____ doses Weekly x _____ doses
 - Monthly x _____ doses Other: _____
- Flush with 0.9% sodium chloride at the completion of infusion
- Patient required to stay for 30 - min observation period
- Total doses:** 1 yr Other

Provider Name (Print) _____ Provider Signature _____ Date _____

ORDERING PROVIDER

Signature X _____ Date _____

Provider _____ Phone _____ Fax _____