TN 100 Covey Drive Suite 307 Franklin, TN 37067





(Crysvita)

Provider_

Burosumab-twza

Infusion orders	Dafe:	
PATIENT INFORMATION		
Name:	DOB: SEX: M	□ F □
ICD-10 code (required):	ICD-10 description:	
□NKDA Allergies:	Weight	lbs/kg:
REFERRAL STATUS		
□New Referral □Referral Renewal □Medication/Order Change □Benefits Verification Only □Discontinuation Order		
PHYSICIAN INFORMATION		
Referral Coordinator Name:	Referral Coordinator Email:	
Ordering Provider:	Provider NPI:	
Referring Practice Name:	Phone: Fax:	
Practice Address:	City: State: Zip Coo	de:
DIAGNOSIS (and ICD 10 code) □ XLH: (familial hypophosphatemia) ICD-10 Code: E83.31 □ TIO: other adult osteomalacia ICD-10 Code: M83.8 □ Other disorders of phosphorus metabolism ICD-10 Code: E83.39	Burosumab-twza ORDERS Indication Pediatric XLH (6 months and older) Adult XLH Pediatric TIO 2 years and older Adult TIO *Adult TIO	
NOTE List Tried & Failed Therapies, including duration of treatment: 1) 2) **Referring physician is responsible for monitoring and reviewing the following labs prior to treatment: • Fasting phosphorus level prior to each dose for first 3 doses and administer only if below ULN • Fasting phosphorus level 2-4 weeks after dose modifications If dose adjustments are needed, new order must be sent by provider based on PI dose calculations	Medication(check one) □ Crysvita less than 10 kg □ Crysvita greater than 10 kg □ Crysvita Dosing □ 1 mg/kg SQ rounded to the nearest 1 mg max 90 mg □ 0.8 mg/kg SQ rounded to the nearest 10 mg max 90 mg □ 1 mg/kg SQ rounded to the nearest 10 mg max 90 mg □ 0.4 mg/kg SQ rounded to the nearest 10 mg □ 0.5 mg/kg not to exceed 180 mg □ 0.5 mg/kg not to exceed 180mg □mg/kg (dose may be increased up to 2mg/kg not to exceed 180mg administered every 2weeks) Frequency □ Every 2 weeks □ Every 4 weeks □ Every 4 weeks □ Everyweeks □ Everyweeks □ Everyweeks □ Everyweeks	
REQUIRED DOCUMENTATION: □ This signed order form by the provider □ Patient demographics AND insurance information □ Clinical/Progress notes supporting primary diagnosis □ Documentation that pt has stopped phos meds and Vit D □ Fasting serum phosphorus concentration should be below the reference range for age prior to initiation of treatment		
ORDERING PROVIDER Signature X	Date	

_____ Phone _____ Fax _