*TN* 100 Covey Drive Suite 307 Franklin, TN 37067





Idursulfase (Elaprase)

radioanase (Liapiase)	ъ.
Provider Order Form	Date:
riovidei Oidei roilli	

Provider Order Form			
PATIENT INFORMATION			
Name:	DOB:	SEX: M □ F □	
ICD-10 code (required):	ICD-10 description:		
□NKDA Allergies:		Weight lbs/kg:	
REFERRAL STATUS			
□New Referral □Referral Renewal □Medication/Order Change □Benefits Verification Only □Discontinuation Order			
PHYSICIAN	INFORMATION		
Referral Coordinator Name:	Referral Coordinator Email:		
Ordering Provider:	Provider NPI:		
Referring Practice Name:	Phone: Fax:		
Practice Address:	City: State:	Zip Code:	
LABORATORY ORDERS  CBC  at each dose every	THERAPY ADMINISTRATION  ☐ Idursulfase (Elaprase) in 100ml 0.9% sodium chloride, intravenous infusion  ■ Dose: 0.5mg/kg  ■ Route: ☐ intravenous  ■ Frequency: once every week  The total volume of infusion should be administered over aperiod of 3 hours, which may be gradually reduced to 1 hourif no hypersensitivity reactions are observed.  ☐ Infuse with a low-protein-binding 0.2 micrometer (OEºm) in-line filter.  ☐ Flush with 0.9% sodium chloride at infusion completion  ☐ Patient is required to stay for 30-minute observation period  ☐ Refills: ☐ Zero / ☐ for 12 months / ☐		
NOTES/ADDITIONAL COMMENTS:  ORDERING PROVIDER			
Signature X	Date		
Drovidor	Dhono Fay		